

Radar Observations of Storms that Produce Sprites or Lightning with Large Charge Moment Changes

Timothy J. Lang, Walter A. Lyons, Steven A. Rutledge, Brenda Dolan, Steven A. Cummer, Paul Krehbiel, William Rison, and Lizxandra Flores-Rivera

AMS Radar Conference
Poster #172



FMA Research, Inc.



Duke
UNIVERSITY


NEW MEXICO TECH
SCIENCE • ENGINEERING • RESEARCH • UNIVERSITY



1. Introduction


What we think we know:

- Charge moment change (CMC) in cloud-to-ground lightning (CGs): $CMC = Q \times Z$ (charge times height)
- The bigger the CMC, the more likely a sprite ($> 100\text{-}300\text{ C km}$ ideal)
- Positive sprites: +CGs tapping broad stratiform charge layers in mesoscale convective systems (MCSs)



Sprite-parent +CG on 19 August 2009 (triangle). Black Dots - VHF sources from OKLMA (Red during sprite). Diamond - Initiation.

- Negative sprites: powerful convective -CGs, also in MCSs



Large-CMC -CG (973 C km) on 26 March 2011 (triangle). Black/Red Dots - VHF sources from NALMA. Diamond - Initiation.

What we'd like to learn about:


- Microphysics and kinematics of storms that produce sprites or large CMCs
- Unusual cases: sub-MCS storms, sprite-parent +CGs in convection

2. Data

- CSU-CHILL Polarimetric Doppler Radar
- CSU-Pawnee Doppler Radar
- NOAA NMQ Radar Mosaics
- Colorado/Oklahoma LMAs
- Duke CMC Network (CMCN)
- Vaisala NLDN




Contact Info: Timothy Lang, NASA MSFC (ZP11), Huntsville, AL 35894; (256) 961-7861, timothy.j.lang@nasa.gov
Funding for this research has come from NASA and DARPA

3. Sprite-Parent +CGs in Smaller Storms





Diamond = Initiation
Big X = Sprite +CG
Dots = LMA Sources
(red = CG to CG+100 ms)
Small Xs = +CGs
Small triangles = -CGs
Gridline spacing = 0.2 deg



8 June 2012 was a severe hailstorm. SP+CGs in stratiform regions initiated in convection. Charge advection allowed weak convection to initiate too.



30 March 2012 OK supercell produced two sprites in ~15 min. Stratiform SP+CGs initiated in convection.




4. Sprite-Parent +CGs in Convection



Some SP+CGs in 8 June 2012 occurred in convection. Evidence of shear causing tilted dipole.

The 25 June 2012 storm produced SP+CGs within convection. It featured inverted charge. The parent flashes discharged mid-level anvil.




Hail shaft melting to heavy rain

Positive phase shift (horizontal alignment)


Negative phase shift (vertical alignment)

Rain

5. Future Work - 31 May/1 June 2013 (El Reno)



Early - 2300 hour Tornado Supercell Large +CMCs in convection



Late - 0700 hour Huge MCS/MCC Large -CMCs in convection, +CMCs in stratiform

+CMCs: Xs
-CMCs: Triangles
($>100\text{ C km}$)
Circle: OKLMA
200-km range

6. Conclusions

- Sprite-parent/large-CMC +CGs need adjacent convective & stratiform/anvil regions with charge pathway between them.
- Convective sprite-parent/large-CMC +CGs associated with mid-level positive charge or tilted dipole.
- El Reno case an archetypal example of both conventional and unconventional large-CMC lightning.